



Conversation Guide

Learning if EPCLUSA is right for you starts with speaking to a Hep C Specialist. EPCLUSA is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C (Hep C) genotype 1, 2, 3, 4, 5, or 6 infection with or without cirrhosis (compensated). In those with advanced cirrhosis (decompensated), EPCLUSA is used with ribavirin.

Use this guide to help your Hep C Specialist understand you're ready to put Hep C behind you.

1

Be prepared to tell your Hep C Specialist about your medical history, including your history with Hep C and any previous Hep C treatment.

A. How long have you had Hep C?

- 0-10 years 11-20 years 21+ years Not sure

B. Have you previously been treated for Hep C?

- Yes No

C. Do you have cirrhosis?

- Yes No Not sure

D. List anything else that may be affecting your health right now, such as liver problems other than Hep C; kidney problems or if you are on dialysis; or any other medical condition.

Important Safety Information

What is the most important information I should know about EPCLUSA?

EPCLUSA can cause serious side effects, including:

- **Hepatitis B virus reactivation:** Before starting EPCLUSA treatment, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during and after treatment with EPCLUSA. This may cause serious liver problems including liver failure and death. If you are at risk, your healthcare provider will monitor you during and after taking EPCLUSA.

Please see Important Facts about EPCLUSA including Important Warning on hepatitis B reactivation located on page 4.

2

Be prepared to tell your Hep C Specialist about symptoms you may be experiencing.

Let your specialist know about any of the symptoms below, or any other symptoms you may be experiencing—even if you don't think they seem related to your liver or hepatitis C.

Which of the following symptoms have you had?

- | | | | |
|----------------------------------------------|---------------------------------------------------------------|-------------------------------------------|-------------------------------------|
| <input type="checkbox"/> Fever | <input type="checkbox"/> Tiredness | <input type="checkbox"/> Loss of appetite | <input type="checkbox"/> Nausea |
| <input type="checkbox"/> Vomiting | <input type="checkbox"/> Dark urine | <input type="checkbox"/> Abdominal pain | <input type="checkbox"/> Joint pain |
| <input type="checkbox"/> Gray-colored stools | <input type="checkbox"/> Jaundice (yellowing of skin or eyes) | | |
| <input type="checkbox"/> Other _____ | | | |

3

Be prepared to tell your Hep C Specialist about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

EPCLUSA and certain other medicines may affect each other, or may cause side effects. Tell your healthcare provider if you take any medicines containing tenofovir disoproxil fumarate (ATRIPLA[®], COMPLERA[®], STRIBILD[®], TRUVADA[®], VIREAD[®]).

List all the medicines (prescription and over-the-counter), vitamins, and herbal supplements you take.

Important Safety Information (continued)

What should I tell my healthcare provider before taking EPCLUSA?

- If you have: ever had hepatitis B infection, liver problems other than hepatitis C infection; severe kidney problems or are on dialysis; any other medical condition; or if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed. It is not known if EPCLUSA will harm your unborn baby. If you take EPCLUSA with ribavirin, you should also read the ribavirin Medication Guide for important pregnancy-related information.
- **Tell your healthcare provider and pharmacist about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. EPCLUSA and certain other medicines may affect each other, or may cause side effects. Tell your healthcare provider if you take any medicines containing tenofovir disoproxil fumarate (ATRIPLA[®], COMPLERA[®], STRIBILD[®], TRUVADA[®], VIREAD[®]).

4

Express yourself and be clear. Remember, you are your best advocate. Explain how Hep C makes you feel. Provide as much specific information as possible.

How exactly does Hep C make you feel? How have things changed over time?

Important Safety Information (continued)

What are the possible side effects of EPCLUSA?

- **Serious side effects may also include: Slow heart rate (bradycardia):** EPCLUSA, when taken with amiodarone (Cordarone[®], Nexterone[®], Pacerone[®]), a medicine used to treat certain heart problems, may cause slow heart rate. In some cases slow heart rate has led to death or the need for a pacemaker when amiodarone is taken with medicines containing sofosbuvir and another antiviral medicine. Get medical help right away if you take amiodarone with EPCLUSA and get any of the following symptoms: fainting or near-fainting, dizziness or lightheadedness, not feeling well, weakness, extreme tiredness, shortness of breath, chest pains, confusion, or memory problems.

The most common side effects of EPCLUSA include headache and tiredness. When EPCLUSA is used with ribavirin for patients with advanced cirrhosis, side effects also include anemia, nausea, insomnia, and diarrhea.

Please see Important Facts about EPCLUSA including Important Warning on hepatitis B reactivation located on page 4.



(ep-KLOO-suh)

MOST IMPORTANT INFORMATION ABOUT EPCLUSA

EPCLUSA can cause serious side effects, including:

Hepatitis B virus reactivation: Before starting treatment with EPCLUSA, your healthcare provider will do blood tests to check for hepatitis B infection. If you have ever had hepatitis B, the hepatitis B virus could become active again during or after treatment with EPCLUSA. This may cause serious liver problems including liver failure and death. Your healthcare provider will monitor you if you are at risk for hepatitis B reactivation during and after you stop taking EPCLUSA.

ABOUT EPCLUSA

EPCLUSA is a prescription medicine used to treat adults with chronic (lasting a long time) hepatitis C (Hep C) genotype 1, 2, 3, 4, 5, or 6 infection with or without cirrhosis (decompensated). In those with advanced cirrhosis (decompensated), EPCLUSA is used with ribavirin.

POSSIBLE SIDE EFFECTS OF EPCLUSA

EPCLUSA can cause serious side effects, including:

- Those in the “Most Important Information about EPCLUSA” section.
- **Slow heart rate (bradycardia):** EPCLUSA, when taken with amiodarone (Cordarone®, Nexterone®, Pacerone®), a medicine used to treat certain heart problems, may cause serious side effects, including slow heart rate. In some cases slow heart rate has led to death or the need for a pacemaker when amiodarone is taken with medicines containing sofosbuvir and another antiviral medicine. Get medical help right away if you take amiodarone with EPCLUSA and get any of the following symptoms:
 - fainting or near-fainting
 - dizziness or lightheadedness
 - not feeling well
 - weakness
 - extreme tiredness
 - shortness of breath
 - chest pains
 - confusion
 - memory problems

The most common side effects of EPCLUSA include headache and tiredness. When EPCLUSA is used with ribavirin for those with advanced cirrhosis, they also include anemia, nausea, insomnia, and diarrhea.

These are not all the possible side effects of EPCLUSA. Tell your healthcare provider if you have any new symptoms while taking EPCLUSA.

IMPORTANT FACTS

This is only a brief summary of important information about EPCLUSA and does not replace talking to your healthcare provider about your condition and your treatment.

BEFORE TAKING EPCLUSA

Tell your healthcare provider about all of your medical conditions, including if you:

- have ever had hepatitis B virus infection
- have liver problems other than hepatitis C infection
- have severe kidney problems or you are on dialysis
- have any other medical condition
- are pregnant or breastfeeding, or plan to become pregnant or breastfeed
 - If you take EPCLUSA with ribavirin, you should also read the ribavirin Medication Guide for important pregnancy-related information.

Tell your healthcare provider about all the medicines you take:

- Keep a list that includes all prescription and over-the-counter medicines, vitamins, and herbal supplements, and show it to your healthcare provider.
- Including any medicines containing tenofovir disoproxil fumarate (ATRIPLA®, COMPLERA®, STRIBILD®, TRUVADA®, VIREAD®).

EPCLUSA and certain medicines may affect each other, or cause side effects.

GET MORE INFORMATION

- This is only a brief summary of important information about EPCLUSA. Talk to your healthcare provider or pharmacist to learn more.
- Go to EPCLUSAINFO.com or call 1-844-4-EPCLUSA
- To learn about potential savings, call 1-855-7-MYPATH or go to EPCLUSAINFO.com/support-path



EPCLUSA, the EPCLUSA logo, COMPLERA, STRIBILD, TRUVADA, VIREAD, GILEAD and the GILEAD logo are trademarks of Gilead Sciences, Inc., or its related companies. ATRIPLA is a trademark of Bristol-Myers Squibb & Gilead Sciences, LLC. All other trademarks referenced herein are the property of their respective owners.